arXiv:1411.2935v1 [math.GT] 11 Nov 2014

Higher derivatives of length functions along earthquake deformations

Martin Bridgeman*

November 12, 2014

1 Introduction

Let S be a closed surface of genus $g \geq 2$ and $T(S)$ the associated Teichmüller space of hyperbolic structures on S. Given $\gamma \in \pi_1(S)$, let $L_\gamma : T(S) \to \mathbb{R}$ be the associated length function and $T_\gamma : T(S) \to \mathbb{R}$ the associated trace function. The functions L_γ, T_γ have a simple relation given by

$$T_\gamma = 2\cosh(L_\gamma/2). \tag{1}$$

Let β be the homotopy class of a simple multicurve (i.e. a union of disjoint simple non-trivial closed curves in S) and t_β the vector field on $T(S)$ associated with left twist along the geodesic representative of β (see [4]). In this paper, we describe a formula to calculate the higher order derivatives of the functions L_γ, T_γ along t_β. In particular we will find a formula for

$$t_\beta^k L_\gamma = t_\beta t_\beta \dots t_\beta L_\gamma.$$

The formulae we derive generalize formulae for the first two derivatives due to Kerchoff (1st derivative, see [4]) and Wolpert (1st and 2nd derivatives, see [5, 6]).

Kerckhoff and Wolpert both showed that the first derivative is given by

$$t_\beta L_\gamma = \sum_{p \in \beta' \cap \gamma'} \cos\theta_p \tag{2}$$

where β', γ' are the geodesic representatives of β, γ respectively and θ_p is the angle of intersection at $p \in \beta' \cap \gamma'$. Kerckhoff further generalized the formula for the case when β, γ are measured laminations (see [4]).

*This work was partially supported by grant #266344 from the Simons Foundation

In [6], Wolpert derived this formula for second derivative

$$t_\alpha t_\beta L_\gamma = \sum_{(p,q)\in\beta'\cap\gamma'\times\alpha'\cap\gamma'} \frac{e^{l_{pq}}+e^{l_{qp}}}{2(e^{L_\gamma}-1)}\sin\theta_p\sin\theta_q + \sum_{(r,s)\in\beta'\cap\gamma'\times\beta'\cap\alpha'} \frac{e^{m_{rs}}+e^{m_{sr}}}{2(e^{L_\beta}-1)}\sin\theta_r\sin\theta_s.$$

where l_{xy} is the length along γ between x, y (similarly m_{xy} is the length along β.)

It follows from Wolpert's formula that

$$t_\beta^2 L_\gamma = t_\beta t_\beta L_\gamma = \sum_{p,q\in\beta'\cap\gamma'} \frac{e^{l_{pq}}+e^{l_{qp}}}{2(e^{L_\gamma}-1)}\sin\theta_p\sin\theta_q. \tag{3}$$

Our formula generalizes equations 2, and 3 to higher derivatives. Our approach is to derive a formula for the higher derivatives of T_γ and then use the functional relation in equation 1 to derive the formula for L_γ.

2 Higher Derivative Formula

We take the geodesic representatives of β and γ. We let the geometric intersection number satisfy $i(\beta,\gamma) = n$ and we order the points of intersection $x_1,\ldots,x_n$ by choosing a base point on γ. We let θ_i be the angle of intersection of β,γ at x_i and l_i be the length along γ from x_1 to x_i. This gives us n-tuples $(l_1,\ldots,l_n)$ and $(\theta_1,\ldots,\theta_n)$.

In order to describe the formula for the higher derivatives, we first introduce some more notation.

Given r, we let $P(r)$ be the set of subsets of the set $\{1,\ldots,r\}$. Then for $I\in P(r)$ will be denoted by $I = (i_1,\ldots,i_k)$ where $1\le i_1 < i_2 < \ldots < i_k \le r$. We then define $\hat{I}$ to be the complementary subset. We also let $|I|$ be the cardinality of I.

We define the alternating length L_I for $I = (i_i,\ldots,i_k)$ by

$$L_I = \sum_{j=1}^{k}(-1)^j l_{i_j} = -l_{i_1} + l_{i_2} - l_{i_3} - \ldots + (-1)^k l_{i_k}$$

We further define a signature for $I\in P(r)$. For $I = (i_1,\ldots,i_k)$ we can consider the integers in $\{1,\ldots,r\}$ in the ordered blocks $[1,i_1],[i_1,i_2],\ldots[i_k,r]$. We take the sum of the cardinality of the even ordered blocks. Then

$$s(I) = (i_2-i_1+1)+(i_4-i_3+1)+\ldots(i_k-i_{k-1}+1) \qquad k \text{ even}$$

$$s(I) = (i_2-i_1+1)+(i_4-i_3+1)+\ldots(r-i_k+1) \qquad k \text{ odd}$$

For $(\theta_1,\ldots,\theta_n)$ we also define

$$\cos(\theta_I) = \prod_{j=1}^{k}\cos(\theta_{i_j}) = \cos(\theta_{i_1})\cos(\theta_{i_2})\ldots\cos(\theta_{i_k})$$

and similarly define $f(\theta_I)$ for f a trigonometric functions.

We let $u_j = l_j + i\theta_j$. The function F_r is given by

$$F_r(u_1, \ldots, u_r, L) = \sum_{I \in P(r), |I| \text{ even}} (-1)^{s(I)} \sin(\theta_I) \cos(\theta_{\hat{I}}) \left(e^{L/2 - L_I} + (-1)^r e^{L_I - L/2} \right).$$

or equivalently

$$F_r(u_1, \ldots, u_r, L) = \sum_{I \in P(r), |I| \text{ even}} (-1)^{s(I)} 2 \sin(\theta_I) \cos(\theta_{\hat{I}}) \cosh(L/2 - L_I)$$

for r even and

$$F_r(u_1, \ldots, u_r, L) = \sum_{I \in P(r), |I| \text{ even}} (-1)^{s(I)} 2 \sin(\theta_I) \cos(\theta_{\hat{I}}) \sinh(L/2 - L_I)$$

for r odd.

We let $C(n, r)$ be the set of the subsets of size r of the set $\{1, 2, \ldots, n\}$. It is given by

$$C(n, r) = \{I = (i_1, i_2, \ldots, i_r) \mid 1 \leq i_1 < i_2 < \ldots < i_r \leq n\}$$

Given $m \in \mathbb{N}$, we let $[m]$ be the parity of m, i.e. $[m] = 0$ if m is even, and $[m] = 1$ if m is odd.

Theorem 1 *Let β be a homotopy class of a simple closed multicurve and γ a homotopy class of non-trivial closed curve. Let the geometric intersection number $i(\beta, \gamma) = n$. Then*

$$t_\beta^k T_\gamma = \frac{1}{2^k} \sum_{\substack{r=0 \\ [r]=[k]}}^{k} B_{n,k,r} \sum_{I \in C(n,r)} F_r(u_{i_1}, \ldots, u_{i_r}, L_\gamma)$$

where $B_{n,k,r}$ are constants described below.

The first two equations correspond to formulae 2 and 3 for the derivatives of length. We use the above, to derive the next case as an example.

Third Derivative: We use the above formula to calculate the formula for the third derivative.

$$t_\beta^3 T_\gamma = \frac{1}{8}((6n-4) \sinh(L_\gamma/2) \sum_{i=1}^{n} \cos(\theta_i) + 12(\sum_{i<j<k} \sinh(L_\gamma/2) \cos(\theta_i) \cos(\theta_j) \cos(\theta_k)$$

$$+ \sinh(L_\gamma/2 - l_{ij}) \sin(\theta_i) \sin(\theta_j) \cos(\theta_k) - \sinh(L_\gamma/2 - l_{ik}) \sin(\theta_i) \cos(\theta_j) \sin(\theta_k)$$

$$+ \sinh(L_\gamma/2 - l_{jk}) \cos(\theta_i) \sin(\theta_j) \sin(\theta_k))$$

2.1 Constants $B_{n,k,r}$

We let $P(k,n)$ be the collection of partitions of k into n ordered nonnegative integers, i.e.

$$P(k,n) = \left\{ p = (p_1, p_2, \ldots, p_n) \in {\mathbb{N}_0}^n \;\middle|\; \sum_{i=1}^{n} p_i = k \right\}$$

For $p \in P(k,n)$, we define $[p] = ([p_1], \ldots, [p_n])$ where $[n]$ is the parity of n. We let $|p| = [p_1] + \ldots [p_n]$. Then $[p]$ is an n-tuple of 0's and 1's with exactly $|p|$ 1's.

Given $p \in P(k,n)$ we define $B(p)$ as a sum of multnomials given by

$$B(p) = \sum_{q \in P(k,n), [q]=[p]} \left(\begin{array}{c} k \\ q \end{array} \right).$$

It is easy to see that $B(p)$ only depends on n, k and $r = |p|$. We therefore define

$$B_{n,k,r} = B(p) \qquad \text{for some } p \text{ with } |p| = r$$

In particular if we let $p_r = (1, 1, \ldots, 1, 0, \ldots, 0) \in P(k,n)$, of r 1's followed by $(n-r)$ 0's, we have

$$B_{n,k,r} = \sum_{p \in P(k,n), [p]=[p_r]} \left(\begin{array}{c} k \\ p \end{array} \right).$$

A simple calculation gives

$$B_{n,k,k} = \left(\begin{array}{c} k \\ p_k \end{array} \right) = \left(\begin{array}{c} k \\ 1, 1, 1 \ldots, 0, 0, \ldots 0 \end{array} \right) = k!$$

3 Twist Deformation

We consider $T(S)$ as the fuchsian locus of the associated quasifuchsian space $QF(S)$. Let $X \in T(S)$ and $X = \mathbb{H}^2/\Gamma$ where Γ is a subgroup of $PSL(2,\mathbb{C})$ acting on upper half space $\mathbb{H}^3 = \{(u,v,w) \in \mathbb{R}^3 \mid w > 0\}$ fixing the hyperbolic plane $\mathbb{H}^2 = \{(u,0,w) | w > 0\}$. Let Γ_z be the subgroup of $PSL(2,\mathbb{C})$ obtained by complex shear-bend along β by amount $z = s + it$, i.e. left shear by amount s followed by bend of t. Then for small z, $X_z = \mathbb{H}^3/\Gamma_z \in QF(S)$. In the terminology of Epstin-Marden this is a quake-bend deformation. See II.3 of [3] for details on quake bend deformations and II.3.9 for a detailed discussion of derivatives of length along quakbend deformations.

Let $\gamma \in \Gamma$ be a hyperbolic element and let $\gamma(z) \in \Gamma_z$ be the element of the deformed group corresponding to γ and $L(z)$ the complex translation length of $\gamma(z)$. To see how γ is deformed, by conjugating, we assume that γ has axis the geodesic g with endpoints $0, \infty \in \hat{\mathbb{C}}$ and is given by

$$\gamma = \left(\begin{array}{cc} \lambda & 0 \\ 0 & 1/\lambda \end{array} \right) \text{ with } \lambda = e^{L/2} \text{ where } L > 0 \text{ is the translation length of } \gamma.$$

We consider the lifts of β which intersect the axis g of γ and normalize to have a lift of β labelled β_1 which intersects axis g at height 1. We enumerate all other lifts by the order of the height of their intersection point with g starting with the intersection point of β_1. Let n be such that $\gamma\beta_1 = \beta_{n+1}$. Let $R_i(z)$ be the Möbius transformation corresponding to a complex bend about β_i of z. Then under the complex bend about β, $\gamma(z)$ given by

$$\gamma(z) = R_1(z)R_2(z)\dots R_n(z)\gamma.$$

A similar description of the deformation of an element in the punctured surface case can be given in terms of shearing coordinates (see [2] for details).

Taking traces we have

$$T(z) = Tr(R_1(z)R_2(z)\dots R_n(z)\gamma) = 2\cosh(L(z)/2).$$

We can find the derivatives of $L(z)$ by differentiating this formula repeatedly. The final formula is obtained by applying symmetry relations on the derivatives and some elementary combinatorics.

We note that both $T(z)$ and $L(z)$ are holomorphic in z. Differentiating in the real direction we have

$$t_\beta^k L_\gamma = \frac{d^k L}{dz^k}(0) = L^{(k)}(0)$$

Also if we let b_β be the vector field on $T(S)$ given by pure bending along β, then we have by analyticity of $L(z)$

$$b_\beta^k L_\gamma = i^k L^{(k)}(0) = (it_\beta)^k L_\gamma.$$

This corresponds to the observation that $b_\beta = J.t_\beta$ where J is the complex structure on $QF(S)$ (see [1]).



Figure 1: Lift of γ

3.1 Derivation of First Two Derivatives

We now calculate the first two derivatives and recover Wolpert's formulae. By the product rule we have

$$T'(0) = \sum_{i=1}^{n} Tr(R_i'(0)\gamma) \qquad T''(0) = \sum_{i=1}^{n} Tr(R_i''(0)\gamma) + 2\sum_{\substack{i,j=1 \\ i<j}}^{n} Tr(R_i'(0)R_j'(0)\gamma) \tag{4}$$

We now describe $R_i(z)$. Let β_i have endpoints $a_i, b_i \in \mathbf{R}$ where $a_i > 0$ and $b_i < 0$. We let λ_i be the height at which β_i intersects g. We orient β_i from a_i to b_i and g from 0 to ∞ and let θ_i be the angle β_i makes with side g with respect to these orientations (see figure 1).

Then

$$\lambda_i = \sqrt{-a_i b_i} \qquad \cos\theta_i = -\left(\frac{a_i + b_i}{a_i - b_i}\right) \qquad \sin\theta_i = \frac{2\sqrt{-a_i b_i}}{a_i - b_i}.$$

As β_1 intersects at height 1, the distance l_i between the intersection points of β_1 and β_i is given by $e^{l_i} = \lambda_i$. Then we let $f_i \in SL(2,\mathbb{R})$ acting on the upper-half space by $f_i(z) = (z - a_i)/(z - b_i)$ and let $S(z) = f_i R_i(z) f_i^{-1}$. Then $S_i(z)$ is the complex translation given by

$$S(z) = \begin{pmatrix} e^{z/2} & 0 \\ 0 & e^{-z/2} \end{pmatrix}.$$

Thus as $R_i(z) = f_i^{-1} S(z) f_i$. Taking derivatives we have $R_i'(0) = f_i^{-1} S'(0) f_i$ and

$$R_i'(0) = \frac{1}{a_i - b_i}\begin{pmatrix} -b_i & a_i \\ -1 & 1 \end{pmatrix}\begin{pmatrix} 1/2 & 0 \\ 0 & -1/2 \end{pmatrix}\begin{pmatrix} 1 & -a_i \\ 1 & -b_i \end{pmatrix} = \frac{1}{2(a_i - b_i)}\begin{pmatrix} -(a_i + b_i) & 2a_i b_i \\ -2 & a_i + b_i \end{pmatrix}$$

Therefore

$$R_i'(0) = \frac{1}{2}\begin{pmatrix} \cos\theta_i & -e^{l_i}\sin\theta_i \\ -e^{-l_i}\sin\theta_i & -\cos\theta_i \end{pmatrix}$$

Also as $S''(0) = \frac{1}{4}I$ we have $R_i''(0) = \frac{1}{4}I$. Using this we have that

$$Tr(R_i'(0)\gamma) = Tr\left(\frac{1}{2}\begin{pmatrix} \cos\theta_i & -e^{l_i}\sin\theta_i \\ -e^{-l_i}\sin\theta_i & -\cos\theta_i \end{pmatrix}\begin{pmatrix} e^{L/2} & 0 \\ 0 & e^{-L/2} \end{pmatrix}\right) = \sinh(L/2)\cos\theta_i$$

$$Tr(R_i'(0)R_j'(0)\gamma) = Tr\left(\frac{1}{4}\begin{pmatrix} \cos\theta_i & -e^{l_i}\sin\theta_i \\ -e^{-l_i}\sin\theta_i & -\cos\theta_i \end{pmatrix}\begin{pmatrix} \cos\theta_j & -e^{l_j}\sin\theta_j \\ -e^{-l_j}\sin\theta_j & -\cos\theta_j \end{pmatrix}\begin{pmatrix} e^{L/2} & 0 \\ 0 & e^{-L/2} \end{pmatrix}\right).$$

$$= \frac{1}{4}\left(\cos\theta_i\cos\theta_j(e^{L/2} + e^{-L/2}) + \sin\theta_i\sin\theta_j\left(e^{L/2+l_i-l_j} + e^{-(L/2+l_i-l_j)}\right)\right) \tag{5}$$

Let l_{ij} be the distance along γ from β_i to β_j with respect to the orientation of γ. Then for $i < j$ we have $l_{ij} = l_j - l_i$, and $l_{ji} = L - l_{ij}$ for $i > j$.

$$Tr(R_i'(0)R_j'(0)\gamma) = \frac{1}{2}(\cos\theta_i\cos\theta_j\cosh(L/2) + \sin\theta_i\sin\theta_j\cosh(L/2 - l_{ij})). \tag{6}$$

Combining these we obtain the first two derivatives of T_γ.

$$T'(0) = \sinh(L/2)\sum_{i=1}^{n}\cos\theta_i$$

$$T''(0) = \sum_{\substack{i,j=1 \\ i<j}}^{n} (\cos\theta_i \cos\theta_j \cosh(L/2) + \sin\theta_i \sin\theta_j \cosh(L/2 - l_{ij})) + \frac{n\cosh(L/2)}{2}$$

As $T(z) = 2\cosh(L(z)/2)$, then $T'(0) = \sinh(L/2)L'(0)$ giving

$$L'(0) = \sum_{i=1}^{n} \cos\theta_i$$

Also $T''(0) = \frac{1}{2}\cosh(L/2)(L'(0))^2 + \sinh(L/2)L''(0)$. Therefore

$$T''(0) = \frac{\cosh(L/2)}{2}\left(n + 2.\sum_{\substack{i,j=1 \\ i<j}}^{n} \cos\theta_i \cos\theta_j\right) + \sum_{\substack{i,j=1 \\ i<j}} \sin\theta_i \sin\theta_j \cosh(L/2 - l_{ij}).$$

We have

$$n + 2.\sum_{\substack{i,j=1 \\ i\neq j}}^{n} \cos\theta_i \cos\theta_j = \left(\sum_{i=1}^{n} \cos\theta_i\right)^2 + \sum_{i=1}^{n} \sin^2\theta_i$$

and

$$T''(0) = \frac{\cosh(L/2)\left((\sum_{i=1}^{n} \cos\theta_i)^2 + \sum_{i=1}^{n} \sin^2\theta_i\right)}{2} + \sum_{\substack{i,j=1 \\ i<j}} \sin\theta_i \sin\theta_j \cosh(L/2 - l_{ij}). \qquad (7)$$

Solving for $L''(0)$ we obtain

$$L''(0) = \sum_{i=1}^{n} \frac{\sin^2\theta_i}{2\tanh(L/2)} + \sum_{\substack{i,j=1 \\ i<j}}^{n} \frac{\sin\theta_i \sin\theta_j \cosh(L/2 - l_{ij})}{\sinh(L/2)}$$

As $l_{ii} = 0$ we can write

$$L''(0) = \sum_{i,j=1}^{n} \frac{e^{l_{ij}-L/2} + e^{L/2-l_{ij}}}{2(e^{L/2} - e^{-L/2})} \sin\theta_i \sin\theta_j = \sum_{i,j=1}^{n} \frac{e^{l_{ij}} + e^{l_{ji}}}{2(e^{L} - 1)} \sin\theta_i \sin\theta_j.$$

The above give the formulae 2 and 3 as described.

4 Higher Derivatives

We now derive the formula for higher derivatives. We have the formula

$$T(z) = Tr(R_1(z)R_2(z)\dots R_n(z)\gamma).$$

We let $P(k,n)$ be the collection of partitions of k into n ordered nonnegative integers, i.e.

$$P(k,n) = \left\{ p = (p_1, p_2, \ldots, p_n) \in {\mathbb{N}_0}^n \;\middle|\; \sum_{i=1}^{n} p_i = k \right\}$$

Then by the product rule, the kth derivative of T at zero is,

$$T^{(k)}(0) = \sum_{p \in P(k,n)} \left(\begin{array}{c} k \\ p \end{array} \right) Tr(R_1^{(p_1)}(0) \ldots R_n^{(p_n)}(0)\gamma)$$

We have from above that $R_i(z) = f_i^{-1} S(z) f_i$ where

$$S(z) = \left(\begin{array}{cc} e^{z/2} & 0 \\ 0 & e^{-z/2} \end{array} \right).$$

As $S^{(2)}(z) = \frac{1}{4} S(z)$, we have for m even

$$R_i^{(m)}(0) = \frac{1}{2^m} I$$

and for m odd we have

$$R_i^{(m)}(0) = \frac{1}{2^{m-1}} R_i'(0) = \frac{1}{2^m} \left(\begin{array}{cc} \cos\theta_i & -e^{l_i} \sin\theta_i \\ -e^{-l_i} \sin\theta_i & -\cos\theta_i \end{array} \right).$$

Let $z = x + iy$ and define

$$A(z) = \left(\begin{array}{cc} \cos y & -e^{x} \sin y \\ -e^{-x} \sin y & -\cos y \end{array} \right).$$

We let $u_j = l_j + i\theta_j$. Then

$$R_j^{(p)}(0) = \left\{ \begin{array}{cc} \frac{1}{2^p} A(u_j) & p \text{ odd} \\ \frac{1}{2^p} I & p \text{ even} \end{array} \right.$$

Therefore

$$T^{(k)}(0) = \frac{1}{2^k} \sum_{p \in P(k,n)} \left(\begin{array}{c} k \\ p \end{array} \right) Tr(A(u_1)^{[p_1]} \ldots A(u_n)^{[p_n]} \gamma)$$

where $[m]$ is the parity of m. We define

$$F_r(z_1, \ldots, z_r, L) = Tr(A(z_1) \ldots A(z_r)\gamma)$$

Therefore gathering terms we have

$$T^{(k)}(0) = \frac{1}{2^k} \sum_{r=0}^{k} B_{n,k,r} \sum_{1 \le i_1 < \ldots < i_r \le n} F_r(u_{i_1}, \ldots, u_{i_r}, L)$$

where $B_{n,k,r}$ are the coefficients described above. We note that we only get non-zero terms for $[r] = [k]$, so we have $B_{n,k,r} = 0$ for $[k] \neq [r]$.

We define the function

$$G_r(u_1,\dots,u_n,L)=\sum_{I\in C(n,r)} F_r(u_{i_1},\dots,u_{i_r},L).$$

Then G_r is symmetric in $(u_1,\dots,u_n)$ and we have

$$t_\beta^k T_\gamma=\frac{1}{2^k}\sum_{\substack{r=0\\ [r]=[k]}}^{k} B_{n,k,r}G_r(u_1,\dots,u_n,L_\gamma)$$

4.1 Function F_r

We now calculate the formula for F_r.

Lemma 1 *The function F_r is given by*

$$F_r(u_1,\dots,u_r,L)=\sum_{I\in P(r),|I|\ even}(-1)^{s(I)}\sin(\theta_I)\cos(\theta_{\hat{I}})\left(e^{L/2-L_I}+(-1)^r e^{L_I-L/2}\right).$$

or equivalently

$$F_r(u_1,\dots,u_r,L)=\sum_{I\in P(r),|I|\ even}(-1)^{s(I)}2\sin(\theta_I)\cos(\theta_{\hat{I}})\cosh(L/2-L_I)$$

for r even and

$$F_r(u_1,\dots,u_r,L)=\sum_{I\in P(r),|I|\ even}(-1)^{s(I)}2\sin(\theta_I)\cos(\theta_{\hat{I}})\sinh(L/2-L_I)$$

for r odd.

Proof: We have

$$A(u)=\begin{pmatrix}\cos\theta & -e^{l}\sin\theta\\ -e^{-l}\sin\theta & -\cos\theta\end{pmatrix}.$$

Therefore $F_r(u_1,\dots,u_r,L)=Tr(A(u_1),\dots,A(u_r)\gamma)$ has the form

$$F_r(u_1,\dots,u_r,L)=\sum_{I\in P(r)}a_I\sin(\theta_I)\cos(\theta_{\hat{I}})$$

for some coefficients a_I. Expanding we have

$$\begin{aligned}F_r(u_1,\dots,u_r,L) &= (A(u_1)\dots A(u_r)\gamma)_1^1+(A(u_1)\dots A(u_r)\gamma)_2^2\\ &= e^{L/2}(A(u_1)\dots A(u_r))_1^1+e^{-L/2}(A(u_1)\dots A(u_r))_2^2.\end{aligned}$$

Similarly we have

$$(A(u_1)\dots A(u_r))_j^i=\sum_{I\in P(r)}a_j^i(I)\sin(\theta_I)\cos(\theta_{\hat{I}})$$

and define

$$(A(u_1)\dots A(u_r))^i_j(I) = a^i_j(I)\sin(\theta_I)\cos(\theta_{\hat{I}}).$$

We prove the lemma by induction. Given $I = (i_1, \dots, i_k) \in P(r)$ then $I_j = (i_1, i_2, \dots, i_{j-1}) \in P(i_j)$.

The matrix $A(u)$ has cos terms on the diagonal and sin off diagonal. As $\sin(\theta_{i_k})$ is the last sin term we have in $(A(u_1), \dots, A(u_r))^1_1(I)$ we have

$$(A(u_1)\dots A(u_r))^1_1(I) = (A(u_1)\dots A(u_{i_{k-1}}))^1_2(I_k)(A(u_{i_k})^2_1 A(u_{i_k+1})^1_1 \dots A(u_r)^1_1$$

$$= \cos(\theta_{i_k+1})\dots\cos(\theta_r)\left(-e^{-l_{i_k}}\sin(\theta_{i_k})\right)(A(u_1)\dots A(u_{i_{k-1}}))^1_2(I_k)$$

Now iterating, as the next sin is $\sin(\theta_{i_{k-1}})$ we have

$$(A(u_1)\dots A(u_{i_{k-1}}))^1_2(I_k) = (A(u_1)\dots A(u_{i_{k-2}}))^1_1(I_{k-1})A^1_2(u_{i_{k-1}})A^2_2(u_{i_{k-1}+1})A^2_2(u_{i_{k-1}+2})\dots A^2_2(u_{i_{k-1}}).$$

$$= (A(u_1)\dots A(u_{i_{k-2}}))^1_1(I_{k-1})(-e^{l_{i_{k-1}}}\sin(\theta_{i_{k-1}}))(-\cos(\theta_{i_{k-1}+1}))(-\cos(\theta_{i_{k-1}+2}))\dots(-\cos(\theta_{i_k-1})).$$

Thus we have

$$\frac{(A(u_1), \dots, A(u_r))^1_1(I)}{(A(u_1)\dots A(u_{i_{k-2}}))^1_1(I_{k-1})} =$$

$$(-1)^{i_k - i_{k-1}+1}e^{l_{i_{k-1}}-l_{i_k}}\sin(\theta_{i_{k-1}})\cos(\theta_{i_{k-1}+1})\dots\cos(\theta_{i_k-1})\sin(\theta_{i_k})\cos(\theta_{i_k+1})\cos(\theta_{i_k+2})\dots\cos(\theta_r).$$

As each off-diagonal term switches the index, there must be an even number of off-digonal terms in the trace and therefore $|I|$ is even. Then by induction

$$(A(u_1), \dots, A(u_r)\gamma)^1_1 = (-1)^{s(I)}\sin(\theta_I)\cos(\theta_{\hat{I}})e^{L/2-L_I}$$

where

$$s(I) = (i_2 - i_1 + 1) + (i_4 - i_3 + 1) + \dots (i_k - i_{k-1} + 1)$$

and

$$L_I = \sum_{j=1}^{k}(-1)^j l_{i_j} = -l_{i_1} + l_{i_2} - l_{i_3} - \dots + (-1)^k l_{i_k}$$

Similarly

$$\frac{(A(u_1), \dots, A(u_r))^2_2(I)}{(A(u_1)\dots A(u_{i_{k-2}}))^2_2(I_{k-1})} =$$

$$= (-e^{-l_{i_{k-1}}}\sin(\theta_{i_{k-1}}))(\cos(\theta_{i_{k-1}+1}))(\cos(\theta_{i_{k-1}+2}))\dots(\cos(\theta_{i_k-1}))\left(-e^{l_{i_k}}\sin(\theta_{i_k})\right)(-\cos(\theta_{i_k+1}))\dots(-\cos(\theta_r))$$

$$= (-1)^{r-i_k+2}e^{-l_{i_{k-1}}+l_{i_k}}\sin\theta_{i_{k-1}}\cos(\theta_{i_{k-1}+1})\dots\cos(\theta_{i_k-1})\sin(\theta_{i_k})\cos(\theta_{i_k+1})\cos(\theta_{i_k+2})\dots\cos(\theta_r).$$

Counting negative signs we have $r - s(I) + |I|$ negative signs.

$$(A(u_1), \dots, A(u_r)\gamma)^2_2 = (-1)^{r-s(I)+|I|}\sin(\theta_I)\cos(\theta_{\hat{I}})e^{L_I - L/2}$$

As $|I|$ is even we get

$$(A(u_1), \dots, A(u_r)\gamma)^2_2 = (-1)^{r+s(I)}\sin(\theta_I)\cos(\theta_{\hat{I}})e^{L_I - L/2}$$

giving the result. □

5 Some examples

We have from the calculations in the last section that

$$F_0(L) = 2\cosh(L/2) \qquad F_1(u, L) = 2\sinh(L/2)\cos\theta$$

$$F_2(u_1, u_2, L) = 2(\cos\theta_1\cos\theta_2\cosh(L/2) + \sin\theta_1\sin\theta_2\cosh(L/2 - l_{12}))$$

Calculating F_3 we have

$$F_3(u_1, u_2, u_3, L) = 2\sinh(L/2)\cos(\theta_1)\cos(\theta_2)\cos(\theta_3) + 2\sinh(L/2 - l_{12})\sin(\theta_1)\sin(\theta_2)\cos(\theta_3)$$

$$-2\sinh(L/2 - l_{13})\sin(\theta_1)\cos(\theta_2)\sin(\theta_3) + 2\sinh(L/2 - l_{23})\cos(\theta_1)\sin(\theta_2)\sin(\theta_3)$$

Therefore we have

$$G_0(L) = 2\cosh(L/2) \qquad G_1(u_1, \ldots, u_n, L) = 2\sinh(L/2)\sum_{i=1}^{n}\cos\theta_i$$

$$G_2(u_1, \ldots, u_n, L) = 2\sum_{\substack{i,j=1 \\ i<j}}^{n}(\cos\theta_i\cos\theta_j\cosh(L/2) + \sin\theta_i\sin\theta_j\cosh(L/2 - l_{ij}))$$

$$G_3(u_1, \ldots, u_n, L) = 2\sum_{i<j<k}(\sinh(L/2)\cos(\theta_i)\cos(\theta_j)\cos(\theta_k) + \sinh(L/2 - l_{ij})\sin(\theta_i)\sin(\theta_j)\cos(\theta_k)$$

$$-\sinh(L/2 - l_{ik})\sin(\theta_i)\cos(\theta_j)\sin(\theta_k) + \sinh(L/2 - l_{jk})\cos(\theta_i)\sin(\theta_j)\sin(\theta_k))$$

As the functions G_r do not depend on k, once we've calculated all derivatives less than k, we only need calculate G_k to find the kth derivative.

For $k = 3$ we have

$$t_\beta^3 T_\gamma = \frac{1}{8}\left(B_{n,3,1}G_1(u_1, \ldots, u_n, L_\gamma) + B_{n,3,3}G_3(u_1, \ldots, u_n, L_\gamma)\right)$$

$$B_{n,3,3} = 3! = 6 \qquad B_{n,3,1} = (n-1)\begin{pmatrix} 3 \\ 1,2 \end{pmatrix} + \begin{pmatrix} 3 \\ 3 \end{pmatrix} = 3(n-1) + 1 = 3n - 2$$

$$t_\beta^3 T_\gamma = \frac{1}{8}((6n-4)\sinh(L_\gamma/2)\sum_{i=1}^{n}\cos(\theta_i) + 12\left(\sum_{i<j<k}\sinh(L_\gamma/2)\cos(\theta_i)\cos(\theta_j)\cos(\theta_k)\right.$$

$$+\sinh(L_\gamma/2 - l_{ij})\sin(\theta_i)\sin(\theta_j)\cos(\theta_k) - \sinh(L_\gamma/2 - l_{ik})\sin(\theta_i)\cos(\theta_j)\sin(\theta_k)$$

$$\left.+\sinh(L_\gamma/2 - l_{jk})\cos(\theta_i)\sin(\theta_j)\sin(\theta_k)\right)$$

References

[1] F. Bonahon, "Shearing hyperbolic surfaces, bending pleated surfaces and Thurston's symplectic form," *Ann. Fac. Sci. Toulouse Math.* (6) **5** (1996), pp. 233- 297.

[2] L. Chekhov, Lecture Notes on Quantum Teichmuller Theory, preprint 2007, arXiv:0710.2051

[3] D.B.A. Epstein and A. Marden, "Convex hulls in hyperbolic space, a theorem of Sullivan, and measured pleated surfaces,"in *Fundamentals of Hyperbolic Geometry: Selected Expositions*, R.D. Canary, D.B.A. Epstein, A. Marden, eds., LMS Lecture Notes 328, 2005.

[4] S. Kerckhoff "The Nielsen realization problem, " *Annals of Math.* **117 (2)** (1983), no. 2, pp. 235–265.

[5] S. Wolpert, "An elementary formula for the Fenchel-Nielsen twist", *Comm. Math. Helv.*, **56**, 132-135 (1981)

[6] S. Wolpert, "Thurston's Riemannian metric for Teichmüller space," *J. Diff. Geom.,* **23** (1986), pp. 143–174.